ΑD	)	

Award Number: DAMD17-99-1-9155

TITLE: 4-Aminobiphenyl (4-ABP)-DNA Damage in Breast Tissue and Relationship to p53 Mutation and Polymorphisms of Metabolizing Genes

PRINCIPAL INVESTIGATOR: Nancy J. Niguidula

CONTRACTING ORGANIZATION: Columbia University

New York, New York 10032

REPORT DATE: September 2001

TYPE OF REPORT: Annual Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;

Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

# REPORT DOCUMENTATION PAGE

Form Approved OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching esting data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blan		3. REPORT TYPE AND DATES CO			
4. TITLE AND SUBTITLE	September 2001	Annual Summary (15 Aug	I UU - 14 Aug U1) IG NUMBERS		
4-Aminobiphenyl (4-ABF	99-1-9155				
Relationship to p53 Mutation and Polymorphisms of					
Metabolizing Genes					
Metabolizing denes					
6. AUTHOR(S)					
Nancy J. Niguidula					
7. PERFORMING ORGANIZATION I	8. PERFOR	MING ORGANIZATION			
Columbia University		NUMBER			
New York, New York 10					
E-Mail: njn‡@cu.edu					
O CHONCODING / MCAUTORING	OCENIOV NAMETON AND ADDRESSE	6) 40 0001	CODING / MONITODING		
9. SPONSOKING / MONITORING A	AGENCY NAME(S) AND ADDRESS(E		SORING / MONITORING CY REPORT NUMBER		
U.S. Army Medical Research an	d Materiel Command	Agen	or the our monden		
Fort Detrick, Maryland 21702-5					
11. SUPPLEMENTARY NOTES					
12a. DISTRIBUTION / AVAILABILIT			12b. DISTRIBUTION CODE		
	Y STATEMENT lease; Distribution Un	limited	12b. DISTRIBUTION CODE		
		limited	12b. DISTRIBUTION CODE		
		limited	12b. DISTRIBUTION CODE		
Approved for Public Re	lease; Distribution Un	limited	12b. DISTRIBUTION CODE		
	lease; Distribution Un	limited	12b. DISTRIBUTION CODE		
Approved for Public Re	elease; Distribution Uni				
Approved for Public Re  13. ABSTRACT (Maximum 200 We  The proposed research proj	elease; Distribution Uni	between environmental exposures	and genetic susceptibility		
Approved for Public Re  13. ABSTRACT (Maximum 200 We  The proposed research proj factors. This project determine	elease; Distribution Universely Distribution Universel		and genetic susceptibility		
Approved for Public Re  13. ABSTRACT (Maximum 200 We  The proposed research proj factors. This project determined by PCR-RFI CYP1A2, NAT2, NAT1 and	elease; Distribution Universely.  Determined the level of 4-aminobiphenyl( LP. The association between 4-A;  GSTP1 were examined. Another	between environmental exposures ABP)-DNA adduct levels with imm BP-DNA adduct levels and the car aim of this project is to determine:	and genetic susceptibility nunhistochemical staining in cinogen metabolizing genes f exposure to 4-ABP and its		
13. ABSTRACT (Maximum 200 Weather The proposed research projectors. This project determined by PCR-RFI CYP1A2, NAT2, NAT1 and metabolites can cause mutati	elease; Distribution United the level of 4-aminobiphenyl(.P. The association between 4-A; GSTP1 were examined. Another tons in the p53 gene. The elucid	between environmental exposures ABP)-DNA adduct levels with imm BP-DNA adduct levels and the car	and genetic susceptibility nunhistochemical staining in cinogen metabolizing genes f exposure to 4-ABP and its		
Approved for Public Re  13. ABSTRACT (Maximum 200 We  The proposed research proj factors. This project determined by PCR-RFI CYP142, NAT2, NAT1 and	elease; Distribution United the level of 4-aminobiphenyl(.P. The association between 4-A; GSTP1 were examined. Another tons in the p53 gene. The elucid	between environmental exposures ABP)-DNA adduct levels with imm BP-DNA adduct levels and the car aim of this project is to determine:	and genetic susceptibility nunhistochemical staining in cinogen metabolizing genes f exposure to 4-ABP and its		
Approved for Public Re  13. ABSTRACT (Maximum 200 We  The proposed research proj factors. This project determined by PCR-RFI CYP1A2, NAT2, NAT1 and metabolites can cause mutati	elease; Distribution United the level of 4-aminobiphenyl(.P. The association between 4-A; GSTP1 were examined. Another tons in the p53 gene. The elucid	between environmental exposures ABP)-DNA adduct levels with imm BP-DNA adduct levels and the car aim of this project is to determine:	and genetic susceptibility nunhistochemical staining in cinogen metabolizing genes f exposure to 4-ABP and its		
Approved for Public Re  13. ABSTRACT (Maximum 200 We  The proposed research proj factors. This project determined by PCR-RFI CYP1A2, NAT2, NAT1 and metabolites can cause mutati	elease; Distribution United the level of 4-aminobiphenyl(.P. The association between 4-A; GSTP1 were examined. Another tons in the p53 gene. The elucid	between environmental exposures ABP)-DNA adduct levels with imm BP-DNA adduct levels and the car aim of this project is to determine:	and genetic susceptibility nunhistochemical staining in cinogen metabolizing genes f exposure to 4-ABP and its		
Approved for Public Re  13. ABSTRACT (Maximum 200 We  The proposed research proj factors. This project determined by PCR-RFI CYP1A2, NAT2, NAT1 and metabolites can cause mutati	elease; Distribution United the level of 4-aminobiphenyl(.P. The association between 4-A; GSTP1 were examined. Another tons in the p53 gene. The elucid	between environmental exposures ABP)-DNA adduct levels with imm BP-DNA adduct levels and the car aim of this project is to determine:	and genetic susceptibility nunhistochemical staining in cinogen metabolizing genes f exposure to 4-ABP and its		
Approved for Public Re  13. ABSTRACT (Maximum 200 We  The proposed research proj factors. This project determined by PCR-RFI CYP1A2, NAT2, NAT1 and metabolites can cause mutati	elease; Distribution United the level of 4-aminobiphenyl(.P. The association between 4-A; GSTP1 were examined. Another tons in the p53 gene. The elucid	between environmental exposures ABP)-DNA adduct levels with imm BP-DNA adduct levels and the car aim of this project is to determine:	and genetic susceptibility nunhistochemical staining in cinogen metabolizing genes f exposure to 4-ABP and its		
13. ABSTRACT (Maximum 200 Weather The proposed research projectors. This project determined by PCR-RFI CYP1A2, NAT2, NAT1 and metabolites can cause mutati	elease; Distribution United the level of 4-aminobiphenyl(.P. The association between 4-A; GSTP1 were examined. Another tons in the p53 gene. The elucid	between environmental exposures ABP)-DNA adduct levels with imm BP-DNA adduct levels and the car aim of this project is to determine:	and genetic susceptibility nunhistochemical staining in cinogen metabolizing genes f exposure to 4-ABP and its		
13. ABSTRACT (Maximum 200 Weather The proposed research projectors. This project determined by PCR-RFI CYP1A2, NAT2, NAT1 and metabolites can cause mutati	elease; Distribution United the level of 4-aminobiphenyl(.P. The association between 4-A; GSTP1 were examined. Another tons in the p53 gene. The elucid	between environmental exposures ABP)-DNA adduct levels with imm BP-DNA adduct levels and the car aim of this project is to determine:	and genetic susceptibility nunhistochemical staining in cinogen metabolizing genes f exposure to 4-ABP and its		
13. ABSTRACT (Maximum 200 Weather The proposed research projectors. This project determined by PCR-RFI CYP1A2, NAT2, NAT1 and metabolites can cause mutati	elease; Distribution United the level of 4-aminobiphenyl(.P. The association between 4-A; GSTP1 were examined. Another tons in the p53 gene. The elucid	between environmental exposures ABP)-DNA adduct levels with imm BP-DNA adduct levels and the car aim of this project is to determine:	and genetic susceptibility nunhistochemical staining in cinogen metabolizing genes f exposure to 4-ABP and its		
13. ABSTRACT (Maximum 200 Weather The proposed research projectors. This project determined by PCR-RFI CYP1A2, NAT2, NAT1 and metabolites can cause mutati	elease; Distribution United the level of 4-aminobiphenyl(.P. The association between 4-A; GSTP1 were examined. Another tons in the p53 gene. The elucid	between environmental exposures ABP)-DNA adduct levels with imm BP-DNA adduct levels and the car aim of this project is to determine:	and genetic susceptibility nunhistochemical staining in cinogen metabolizing genes f exposure to 4-ABP and its		
13. ABSTRACT (Maximum 200 Weather The proposed research project determined by PCR-RFI CYP1A2, NAT2, NAT1 and metabolites can cause mutatic currently under investigation.	elease; Distribution United the level of 4-aminobiphenyl(.P. The association between 4-A; GSTP1 were examined. Another tons in the p53 gene. The elucid	between environmental exposures ABP)-DNA adduct levels with imm BP-DNA adduct levels and the car aim of this project is to determine:	and genetic susceptibility nunhistochemical staining in cinogen metabolizing genes f exposure to 4-ABP and its p53 for all 150 samples is		
13. ABSTRACT (Maximum 200 Weather The proposed research project determined by PCR-RFI CYP1A2, NAT2, NAT1 and metabolites can cause mutatic currently under investigation.	elease; Distribution University  ect investigates the relationship  ned the level of 4-aminobiphenyl( LP. The association between 4-A GSTP1 were examined. Another  ions in the p53 gene. The elucions	between environmental exposures ABP)-DNA adduct levels with imm BP-DNA adduct levels and the car aim of this project is to determine:	and genetic susceptibility nunhistochemical staining in cinogen metabolizing genes f exposure to 4-ABP and its		
13. ABSTRACT (Maximum 200 Weather The proposed research project determined by PCR-RFI CYP1A2, NAT2, NAT1 and metabolites can cause mutatic currently under investigation.	elease; Distribution University  ect investigates the relationship  ned the level of 4-aminobiphenyl( LP. The association between 4-A GSTP1 were examined. Another  ions in the p53 gene. The elucions	between environmental exposures ABP)-DNA adduct levels with imm BP-DNA adduct levels and the car aim of this project is to determine:	and genetic susceptibility nunhistochemical staining in cinogen metabolizing genes f exposure to 4-ABP and its p53 for all 150 samples is		
13. ABSTRACT (Maximum 200 Weather The proposed research project determined by PCR-RFI CYP1A2, NAT2, NAT1 and metabolites can cause mutatic currently under investigation.  14. SUBJECT TERMS 4-aminobiphcnyl-DNA according to the proposed research project determined by PCR-RFI CYP1A2, NAT2, NAT1 and metabolites can cause mutatic currently under investigation.	ect investigates the relationship ned the level of 4-aminobiphenyl(LP. The association between 4-AGSTP1 were examined. Another ions in the p53 gene. The elucid	between environmental exposures ABP)-DNA adduct levels with imm BP-DNA adduct levels and the car aim of this project is to determine lation of the mutational spectra of	and genetic susceptibility nunhistochemical staining in cinogen metabolizing genes f exposure to 4-ABP and its p53 for all 150 samples is  15. NUMBER OF PAGES  3 16. PRICE CODE		
13. ABSTRACT (Maximum 200 Weather The proposed research project determined by PCR-RFI CYP1A2, NAT2, NAT1 and metabolites can cause mutatic currently under investigation.	elease; Distribution University  ect investigates the relationship  ned the level of 4-aminobiphenyl( LP. The association between 4-A GSTP1 were examined. Another  ions in the p53 gene. The elucions	between environmental exposures ABP)-DNA adduct levels with imm BP-DNA adduct levels and the car aim of this project is to determine:	and genetic susceptibility nunhistochemical staining in cinogen metabolizing genes f exposure to 4-ABP and its p53 for all 150 samples is		

#### 4-ABP-DNA adducts

The immunohistochemical analysis of 4-ABP-DNA adducts in breast tumor tissue was completed last year. This year the analysis of normal adjacent tissue was determined and a more extensive statistical analysis was completed. Analysis included determining the differences of adduct levels in normal adjacent breast tissue between smokers and nonsmokers. Associations were also determined between 4-ABP-DNA adduct levels and the smoking status variables; pack-years, age at initiation, and smoking intensity.

### Genotyping

The determination of CYP1A2 and NAT2 polymorphisms were completed this year by PCR-RFLP. The Fluorescence Polarization-dye-terminator incorporation assay was abandoned due extremely sensitive nature of the assay and the poor quality of DNA acquired from paraffin-embedded tissue. Genotyping of NAT1 and GSTP1 were also completed this year by PCR-RFLP. The statistical analysis to determine the correlations between genotype and adduct levels in both tumor and normal adjacent tissue has been completed.

## P53 Mutational Spectra

Difficulties arose in obtaining a difference in band migration between the positive and negative controls on the SSCP polyacrylamide gels. This obsolete method has been deserted for the more efficient and comprehensive p53 GeneChip method. The p53 GeneChip probes are commercially available from Affymetrix. The Cancer Core Facility at Columbia University has an Affymetrix Instrument System available to analyze the GeneChips. Complete sequence data for exons 2-11 in p53 will be determined for the 150 samples. Correlations will be made between identified p53 mutations and 4-ABP-DNA adducts.

#### Key Research Accomplishments

- Determined that the 4-ABP-DNA adduct levels in normal adjacent breast tissue is slightly elevated, though not statistically significant, in nonsmokers compared to smokers.
- Determined the 4-ABP-DNA adduct levels in tumor tissue were slightly elevated, though modest and not statistically significant, in smokers with higher exposure in relation to pack-years.
- There were no statistically significant differences in 4-ABP-DNA adduct levels in either tumor or normal adjacent tissue between smokers stratified by age at initiation.
- The relationship between the genotypes for the carcinogen metabolizing enzymes, CYP1A2, NAT2, NAT1 and GSTM1 have in predicting 4-ABP-DNA adduct levels was determined. None of these genes, independently or in combination, had a significant association with adduct levels.
- The analysis of the mutational spectra of p53 is currently under investigation.